

Two-magnon scattering and the spin-phonon interaction beyond the adiabatic approximation

Matthew J. Reilly and A. G. Rojo

Physics Department, The University of Michigan, Ann Arbor, MI 48109-1120

Abstract

We consider a model of Raman scattering for a two-dimensional $S = 1/2$ Heisenberg Anti-Ferromagnet which includes a *dynamical* spin-phonon interaction. We observe a broadening of the line shape due to increased coupling with excited high-energy spin states. Our results are close to a model of random, static exchange interactions, first introduced in this context by Haas *et al.* [J. Appl. Phys. **75**, 6340, (1994)], which, when extended to large numbers of spins, explains experiments in the parent insulating compounds of high- T_c superconductors.

I. INTRODUCTION

Since the discovery of high- T_c superconductors¹, the $S = 1/2$ quantum Heisenberg model has received considerable attention. This is largely due to well-accepted experimental evidence that suggests that these compounds can be described² by a two-dimensional (2D) doped Heisenberg model for spin $S = 1/2$.

One of the experimental techniques used to study the excitations of these systems is Raman scattering. This technique was intensely used in the past for different antiferromagnets of spin $S \geq 1$. For such systems, the main features of the line shape were explained by Parkinson³, who used a spin-wave theory to account for the photon-two-magnon process

involved in the Raman scattering. Very good agreement⁴ was found for K_2NiF_4 , which is well described by a spin $S = 1$ Heisenberg antiferromagnet.

The results of Raman experiments in the parent insulating compounds of high- T_c superconductors, La_2CuO_4 and $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$, show some qualitative differences with the line shape of previously studied antiferromagnets^{5–7}. As in the case of K_2NiF_4 , the line shape is centered at an energy corresponding to a spin-pair excitation. However, in contrast with the case of spin $S = 1$, the linewidth is very broad, and the spectrum has a very long tail that extends beyond the energy of four magnon excitations. Moreover, while the dominant contribution to scattering is in the so-called B_{1g} channel, there is also a significant contribution in the nominally forbidden A_{1g} configuration⁷.

Since for lower spins the quantum fluctuations are larger, some theoretical analysis beyond the mean-field spin-wave approximation has been attempted to explain the broad line shape. Numerical diagonalizations in a 4×4 lattice shows very little structure for the B_{1g} channel: essentially three peaks; a dominant two-magnon peak and two peaks identifiable as four-magnon excitations⁸. This calculation gives vanishing line shape for the A_{1g} channel. Although the structure of the line shape is clearly different from the one observed in the experiment, the first three moments of both lines are in good agreement. The first three moments of the distribution have been calculated in good agreement to experiment by Singh *et al.* using cumulant expansions on a Heisenberg model with diagonal next-nearest-neighbor couplings⁹. Canali and Girvin¹⁰ used the Dyson-Maleev transformation taking into account processes of up to four magnons, and presented convincing evidence that the line shape cannot be explained by quantum fluctuations alone. Raman scattering Hamiltonians with a four-site exchange¹¹ term have been proposed, these increase the relative weight of the four-magnon scattering, but does not explain the broadening of the peaks. This is also consistent with the work of Sugai¹², and Roger and Delrieu¹³. Also, recent studies of spin-pair excitations in a spin $S = 1$ system, NiPS_3 , show a similar linewidth as those observed in the cuprates¹⁴.

From the above considerations, one concludes that it is necessary to invoke an additional

process. It was emphasized by Nori *et al.*¹⁵ that the spin–phonon interaction can be responsible for the broad linewidth, and for the finite cross–section in the otherwise absent A_{1g} channel. They supported their arguments by computing the Raman cross section in finite lattices, using a Heisenberg model with random *static* exchange integrals, and averaging over configurations with equal weight. This calculation is in the spirit of the adiabatic or Born–Oppenheimer approximation¹⁶, that neglects the fluctuations of the phonon field.

Adding phonons to the 2D $S = 1/2$ Heisenberg Hamiltonian (HAF) provides a mechanism¹⁵ for the otherwise forbidden A_{1g} scattering, as well as allowing a coupling between the ground and excited spin states. In a typical HAF, the phonon frequency¹⁵ is about a third of the interaction constant J . The first excited state lies at $3J$, so there is an order of magnitude energy difference between spin and phonon excitations. It was argued¹⁵ that the separation of energy scales justifies the adiabatic phonon approximation.

In the present work we consider the Raman scattering for a Heisenberg model with spin–phonon interaction, including the effects of the phonon dynamics. We solve for the exact ground state of a system in which the vibrational degrees of freedom are allowed a finite number of excitations, an approximation valid for small but finite phonon frequencies. To the best of our knowledge, this is the first calculation of exact diagonalizations for a dynamic spin–phonon system. Some work including phonon dynamics beyond the adiabatic approximation exists in the context of one–dimensional Peierls systems. Fradkin and Hirsh¹⁷ studied the electron–phonon interaction using quantum Monte Carlo simulations, and Proetto and Falicov¹⁸ solved the case of two sites and one phonon.

This paper is organized as follows. In Sec. II we present our Hamiltonian and scattering operator and discuss the rationale for our truncated–phonon model. In Sec. III we present computational results and provide theoretical support for these results. We also present an alternative approach to including phonons and compare it to our model. Sec. IV is devoted to conclusions.

II. MODEL AND PROCEDURE

We study a Heisenberg model with spin–phonon interaction described by the following Hamiltonian:

$$H = \sum_{\langle ij \rangle} \left\{ \left[J - \alpha(a_{ij}^\dagger + a_{ij}) \right] \vec{S}_i \cdot \vec{S}_j + \omega_0 a_{ij}^\dagger a_{ij} \right\} \quad (1)$$

where $\vec{S}_i$ are spin 1/2 operators, $a_{ij}^\dagger(a_{ij})$ is a creation (annihilation) operator for an Einstein phonon of frequency ω_0 , and the symbol “ ij ” refers to a link of a square lattice. In $J_{ij} = J - \delta J_{ij}$ we include the coupling between the phonons and the spin degrees of freedom through a “displacement” operator $\hat{x}_{ij} = \sqrt{\frac{\hbar}{2m\omega_0}}(a_{ij}^\dagger + a_{ij})$. Our parameter α in Eq.(1) is proportional to the spin–phonon coupling constant $\lambda = \alpha\sqrt{2m\omega_0/\hbar}$ relating the change in the exchange integral δJ_{ij} with the displacement: $\delta J_{ij} = \lambda \hat{x}_{ij} = \alpha(a_{ij}^\dagger + a_{ij})$.

Due to computational limitations, we need to restrict ourselves to a small number (6 and 8) of spins. To have these as two dimensional as possible, we placed them in non-periodic ladder type structures as shown in Fig. 1.

We use Einstein phonons to simplify the model, and we consider the highest–energy phonons as they are the closest in energy to the magnon excitations and will thus have the greatest coupling to the magnons. The occupation number of the phonon degrees of freedom at each link is in principle unrestricted; the number of phonons ranges from zero to infinity. This makes the problem intractable from the point of view of exact diagonalizations, since the resulting Hilbert space is infinite. In order to overcome this limitation, we restrict the possible number of phonons at each link by imposing the condition $(a^\dagger)^n = 0$. Computational resources limit us to using $n = 2, 3$. This approximation maps the phonon degree of freedom into a two– or three–state system at each link. The truncation of the phononic occupation states implies that the variations of J_{ij} due to quantum fluctuations are bounded. For example, for $n = 2$, the maximum displacements for a given link is given by the “coherent” states $|\psi_{ij}^\pm\rangle = 2^{-1/2}(1 \pm a_{ij}^\dagger)|0\rangle$, and $\delta J_{ij} = \alpha\langle\psi_{ij}^\pm|(a_{ij}^\dagger + a_{ij})|\psi_{ij}^\pm\rangle = \pm\alpha$. Quantum fluctuations themselves are certainly limited by our truncation, but since they are in general small for

small ω_0 , our approximation will account for the relevant dynamics in the regime $\delta J_{ij}/J < 1$ and $\omega_0/J < 1$.

We are interested in the Raman scattering intensity $I_R(\omega)$, given by:

$$I_R(\omega) = \sum_{\nu} |\langle \nu | \hat{R} | 0 \rangle|^2 \delta(\omega - E_{\nu} + E_0), \quad (2)$$

where $|\nu\rangle$ are the eigenstates and E_{ν} the eigenvalues of H . We compute $I_R(\omega)$ by using the partial fraction expansion method of Gagliano and Balseiro¹⁹. The relevant operator $\hat{R}$ in Eq.(2) depends on the different configurations of the Raman experiment. In general³:

$$\hat{R} = \sum_{\langle ij \rangle} (\vec{E}_{\text{inc}} \cdot \hat{x}_{ij}) (\vec{E}_{\text{scat}} \cdot \hat{x}_{ij}) \vec{S}_i \cdot \vec{S}_j, \quad (3)$$

with $\vec{E}_{\text{inc}}$ and $\vec{E}_{\text{scat}}$ refer to the polarization of the incoming and scattered photon respectively, and $\hat{x}_{ij}$ are unit vectors in the near-neighbor directions of the lattice. For square lattices, two common scattering configurations are the so-called B_{1g} and A_{1g} symmetries. We let the lattice axis lie along the x and y directions, and we define x' and y' the directions along the lattice diagonals ($\hat{x}' = \frac{\hat{y} + \hat{x}}{\sqrt{2}}$ and $\hat{y}' = \frac{\hat{y} - \hat{x}}{\sqrt{2}}$), then B_{1g} corresponds to $\vec{E}_{\text{scat}} \parallel \hat{x}'$ and $\vec{E}_{\text{inc}} \parallel \hat{y}'$, whereas A_{1g} corresponds to both the incident and scattered photon polarized in the same direction: $\vec{E}_{\text{inc}} \parallel \vec{E}_{\text{scat}} \parallel \hat{x}'$.

For the case of the “pure” Heisenberg Hamiltonian (no phonons, or $\alpha = \omega_0 = 0$), the absorption corresponding to the A_{1g} symmetry is zero at any non-zero frequency, since the A_{1g} operator is proportional to the Hamiltonian. We show in the next section how the inclusion of the phonon dynamics gives rise to a finite absorption in this channel, and argue that a spin-phonon interaction accounts for the main features observed in the Raman experiments in the undoped copper oxides. Also, we compare the results obtained in the dynamical model with that of the Heisenberg model with random (static) exchange integrals, which was studied in this context by Nori *et al.*¹⁵

III. RESULTS AND DISCUSSION

In this section we present numerical results for systems of 6 and 8 spins²⁰ corresponding to the geometry shown in Figure 1. We first discuss the A_{1g} symmetry. In this case, and in the absence of lattice distortions, the Raman operator commutes with the Heisenberg Hamiltonian. No line shape is observed in this case: $I_R(\omega) \sim \delta(\omega)$. It was first pointed out by Nori *et al.*¹⁵ that the presence of *static* disorder in the exchange integrals J_{ij} changes the value of the commutator, and can give rise to a finite A_{1g} signal quite similar to experiments. It was argued by Nori *et al.*¹⁵ that the spin–phonon coupling produces disorder in the values of J_{ij} , in the limit where the vibrational motions are much slower than the spin degrees of freedom. Here, we are interested in how this limit is achieved in a system that—up to boundary effects—is translationally invariant, and includes the *dynamics* of the spin–phonon coupling. Consider our Hamiltonian Eq.(1) and the operator $\hat{R}$ for the A_{1g} symmetry. For $\omega_0 \neq 0$, $[H, \hat{R}] \neq 0$ and we expect a finite line shape. In Fig. 2 we show the $I(\omega)$ obtained for $\alpha = 0.1J$ and $\omega_0 = 0.05J$. Some of the qualitative features of the experimental line shape are already present in these finite–size systems: there is a broad line shape of width $\sim 2J$ with a maximum at the two–magnon energy. Note that for our ladder geometries, since most sites have coordination number 3, the two–magnon energy in the Ising limit is located at $2J$, instead of the corresponding $3J$ of the square lattice. Due to the finite size of the lattice, the line shape consists of a series of peaks that are centered at the unperturbed energies, which are indicated by arrows in Fig. 2. These unperturbed energies correspond to the manifold of two–and multi–magnon states. In turn, these “internal” peaks have a finite width that is due to the coupling with the phonons. The energy scale dominating this width is α .

In order to test the validity of the Hilbert space truncation, we have increased the number of allowed excitations per phonon by one, with no qualitative changes in the line shape (see Fig. 3)

It is interesting to consider the following two limiting cases: (a) vanishing spin-phonon

coupling $\alpha \rightarrow 0$ for finite phonon frequency ω_0 , and (b) vanishing phonon frequency $\omega_0 \rightarrow 0$ for finite α . In both limits the A_{1g} line shape vanishes. In (a), the spins are uncoupled from the phonons, and the line shape vanishes because in this limit $[H, \hat{R}] = 0$. Figure 4 illustrates how the line shape vanishes and shifts to lower energies as $\alpha \rightarrow 0$. In limit (b), the phonons are “static”, but remain coupled to the spins, because $\alpha \neq 0$. Since the phonons have no dynamics, the eigenstates can be written in the form

$$|\psi_\nu(\omega_0 = 0)\rangle = |\psi_\nu\{n\}\rangle \otimes |\{n\}\rangle, \quad (4)$$

where $|\{n\}\rangle$ is a static configuration of displacements, in such a way that, for each link ij , one has $(a_{ij}^\dagger + a_{ij})|\{n\}\rangle = \eta_{ij}(\{n\})|\{n\}\rangle$, with $\eta_{ij}(\{n\})$ a c -number. The state $|\psi_\nu\{n\}\rangle$ involves spin degrees of freedom only, and is an eigenstate of the following Hamiltonian

$$H\{n\} = \sum_{\langle ij \rangle} [J - \alpha \eta_{ij}(\{n\})] \vec{S}_i \cdot \vec{S}_j. \quad (5)$$

For $\omega_0 = 0$, the problem is that of an *annealed* configuration of displacements: for each configuration $\{n\}$, the couplings J_{ij} are different, and one has to solve for the spin dynamics in the presence of this distribution of couplings. The complete spectrum is obtained by solving N_p decoupled eigenvalue problems, with N_p being the phononic Hilbert space dimension. For our case of one excitation per link, $N_p = 2^{N_{\text{links}}}$, with N_{links} being the number of links of the lattice. What distinguishes this from a quenched disorder is the fact that the ground state of this problem corresponds to the lowest energy state of the spin system in an *ordered* background of couplings. Since this is also an eigenstate of $\hat{R}$ for the A_{1g} symmetry, the line shape will be zero. In Fig. 5 we show how the line shape vanishes as ω_0 decreases to zero. Note that from this analysis one concludes that, even in the case of $\omega_0 = 0$, the line shape is non-zero at finite temperature. This is due to the presence of thermally excited states, which are eigenstates of a Hamiltonian that does not commute with $\hat{R}$. The right column of Figure 5 shows the rescaled results, which illustrate the line shape approaching a limiting function of ω whose overall amplitude vanishes as $\omega_0 \rightarrow 0$. The main features of this limiting function should be accounted for by a model of static disordered couplings. In previous work

in the context of one-dimensional Peierls systems²¹, it is argued that in the limit of small phonon frequencies, the quantum lattice fluctuations can be modeled by a static, random Gaussian potential with zero mean. A model in which the exchange integrals are taken from a random configuration of couplings was also studied by Nori *et al*¹⁵. We can test this hypothesis in our dynamical model. Results of the comparison between the dynamical and a static disordered system are shown in Fig. 6. Note that both the amplitude and the position of the peaks are in very good mutual agreement. The scattering peaks from the two models have their energies scaled with respect to each other. After they are rescaled, the location and magnitude of the peaks are very close for both models (see Fig. 6). The basis for this rescaling of the horizontal axis is the following. For our dynamic phonon model, the minimum energy state corresponds to the lattice “maximally compressed”: $\eta_{ij} = -\langle x \rangle_0$. The tendency of the system to compress has the effect of renormalizing the interaction coefficient J , since the displacement x_{ij} will tend to be a constant, and therefore $J' = J + \alpha \langle x \rangle_0$. The average displacement will be greater as the possible number of excitations increases; this explains a shift in energy between 1 and 2 excitations as well.

In the model with static disorder, the phonon energy goes to zero, but the fluctuations are still present. In the treatment of Nori *et al.*¹⁵ the argument used is that the phonon energy is small compared to that of the spin excitations, and thus can be neglected. However, we have shown that for ω_0 strictly zero the line shape vanishes. At zero temperature, the disordered model should be compared with the dynamical model at infinitesimal ω_0 . We prove this statement by using perturbation theory in ω_0 . Assume an ordering of the states for $\alpha = \omega_0 = 0$, and let us label those states by the index ν . If $\alpha \ll J$ and $\omega_0 = 0$, each state $|\psi_\nu\rangle$ splits in a manifold $|\psi_\nu\{n\}\rangle$ of almost degenerate states of energies $E_\nu\{n\}$, that correspond to the “disordered” configurations. If we now turn on ω_0 , different states are going to mix, in such a way that the ground state can be written as

$$|\psi_0\rangle \simeq |\psi_0\{n\}_0\rangle + \omega_0 \sum_{\{n\} \neq \{n\}_0} c(\{n\}) |\psi_0\{n\}\rangle \otimes |\{n\}\rangle, \quad (6)$$

where $\{n\}_0$ is the ordered configuration of couplings. A similar expansion can be used for

the excited states. If we use the fact that $A_{1g} \hat{R}$ acts only on the spin degrees of freedom, and that different states $|\{n\}\rangle$ are orthogonal, to lowest order in ω_0 , $I(\omega)$ will be given by

$$I(\omega) \simeq \omega_0^2 \sum_{\{n\}} |c(\{n\})|^2 \sum_{\nu} |\langle \psi_{\nu} \{n\} | \hat{R} | \psi_0 \{n\} \rangle|^2 \delta(\omega - E_{\nu} \{n\} + E_0 \{n\}). \quad (7)$$

We have found computationally a falloff of $I(\omega)$ proportional to ω_0^2 .

Now, if we keep the assumption of $\alpha \ll J$, the states corresponding to each manifold are almost degenerate and we can approximate $c(\{n\}) \sim (M - 1)^{-1/2} \sim M^{-1/2}$, with M the number of configurations $\{n\}$. Therefore, in this limit $I(\omega)$ is given by an average over configurations with *equal* weight.

We now present results for the B_{1g} configuration. In Fig. 7 we compare the B_{1g} scattering from eight $S = 1/2$ Heisenberg spins for three different cases: bare HAF, with phonon interactions, and with static disorder.

We have argued that peaks at higher excitations will eventually merge into one broad peak as the lattice size increases. To show that this is the case, we have calculated the A_{1g} and B_{1g} peaks for a 3×4 non-periodic lattice with static disorder. The results are shown in Fig. 8. These are in good qualitative agreement with experiments, except that the experimental scattering is centered at $\omega \sim 3J$ instead of $2J$ in our case. This is due to finite size effects and to the lattice being non-periodic.

The model with static disorder, when extended to larger numbers of spins than can be obtained with the dynamic phonon model, begins to duplicate experiments. The agreement between the static and dynamic phonon models suggests that the dynamic phonon model will also agree with experiments.

IV. CONCLUSIONS

In this paper we have presented a model of phonon-magnon interaction with a truncated phonon space. We discussed how this model can explain the otherwise forbidden A_{1g} scattering and showed that it does in fact give rise to A_{1g} scattering. We discussed how coupling

to spin excitations leads to a broadening of the line shape. We have proven that phonons can be modeled by static disorder and compared the results obtained by using quenched phonons to our dynamical model. We showed that the amplitude of the A_{1g} scattering should fall off as ω_0^2 . We showed that both the A_{1g} and B_{1g} scattering is broadened. In conclusion, the broadening is due to coupling to spin excitation states through the phonon interaction.

V. ACKNOWLEDGEMENTS

We would like to thank Roberto Merlin and Jim Allen for helpful discussions, and Franco Nori for helpful discussions and a careful reading of this manuscript.

REFERENCES

- ¹ J. G. Bednorz and K. A. Müller, Z. Phys. B **64**, 189 (1986).
- ² E. Manousakis, Rev. Mod. Phys. **63**, 1 (1991).
- ³ J. B. Parkinson, J. Phys. C **2**, 2012 (1969).
- ⁴ P. A. Fleury and H. J. Guggenheim, Phys. Rev. Lett. **24**, 1346 (1970).
- ⁵ S. Sugai, M. Sato, T. Kobayashi, J. Akimitsu, T. Ito, H. Takagi, S. Uchida, S. Hosoya, T. Kajitani, and T. Fukuda, Phys. Rev. B **42**, 1045 (1990).
- ⁶ P. E. Sulewski, P. A. Fleury, K. B. Lyons, S-W. Cheong, and Z. Fisk, Phys. Rev. B **41**, 225 (1990).
- ⁷ K. B. Lyons, P. E. Sulewski, P. A. Fleury, H. L. Carter, A. S. Cooper, G. P. Espinosa, Z. Fisk, and S.-W. Cheong, Phys. Rev. B **39**, 9693 (1989).
- ⁸ E. Gagliano and S. Bacci, Phys. Rev. B **42**, 8772 (1990).
- ⁹ R. R. P. Singh, P. A. Fleury, K. B. Lyons, and P. E. Sulewski, Phys. Rev. Lett. **62**, 2736 (1989).
- ¹⁰ C. M. Canali and S. M. Girvin, Phys. Rev. B **45**, 7127 (1992).
- ¹¹ Y. Honda, Y. Kuramoto, and T. Watanabe, Physica C **185-189**, 1493 (1991).
- ¹² S. Sugai, Solid State Comm. **75**, 795 (1990).
- ¹³ M. Roger and J. M. Delrieu, Synthetic Metals **29**, F673 (1989).
- ¹⁴ S. Rosenblum, A. H. Francis, and R. Merlin, Phys. Rev. B **49**, 4352 (1994).
- ¹⁵ F. Nori, R. Merlin, S. Haas, A. W. Sandvik, and E. Dagotto, Phys. Rev. Lett., **75**, 553 (1995); S. Haas, E. Dagotto, J. Riera, R. Merlin, and F. Nori, J. Appl. Phys. **75**, 6340 (1994).

- ¹⁶ See for example, J. M. Ziman, *Electrons and Phonons* (Oxford University Press, Oxford, 1960), p. 175.
- ¹⁷ E. Fradkin and J. E. Hirsch, Phys. Rev. B **27**, 1680 (1983).
- ¹⁸ C. R. Proetto and L. M. Falicov, Phys. Rev. B **39**, 7545 (1989).
- ¹⁹ E. R. Gagliano and C. A. Balseiro, Phys. Rev. Lett. **59**, 2999 (1987).
- ²⁰ The Hilbert space dimension is $\binom{N}{N/2} n_e^{n_b}$, where N is the number of spins, n_e is the number of allowed phonon states, and n_b is the number of nearest-neighbor pairs. For 6 spins and 2 (3) phonon states in a ladder configuration, this gives a Hilbert dimension of 2560 (43740). For 8 spins, 2 (3) excitations, the dimension becomes 71680 (4133430).
- ²¹ K. Kim, R. H. McKenzie and J. W. Wilkins, Phys. Rev. Lett. **71**, 4015 (1993); R. H. McKenzie and J. W. Wilkins, *ibid* **69**, 1085 (1992); S. A. Brazovskii and I. E. Dzyaloshinskii, Zh. Eksp. Teor. Fiz. **71**, 2338 (1976) [Sov. Phys. JETP **44**, 1233 (1976)].

FIGURES

FIG. 1. Ladder geometries used in our numerical computations. Each dot represents a spin $1/2$. Each solid line connects nearest-neighboring spins, and represents a Heisenberg Anti-Ferromagnetic coupling modulated by an Einstein phonon. There is one phonon per coupling.

FIG. 2. Raman cross section $I(\omega)$ versus ω/J for A_{1g} scattering. The parameters of the dynamical Hamiltonian are $\alpha = 0.1J$, $\omega_0 = 0.05J$, and the number of phonon excitations per link, n , is 2. The vertical scale is arbitrary. The arrows are located at the eigenenergies of the model without phonons corresponding to eigenstates of total spin $S = 0$.

FIG. 3. Comparison between one- and two-phonon (lighter curve) excitations per link, for the Raman line shape $I(\omega)$ in the A_{1g} symmetry. The parameters are $\alpha = 0.1J$, and $\omega_0 = 0.05J$, for a ladder of 6 spins.

FIG. 4. Raman line shape for the A_{1g} symmetry. Here we consider 6 spins (including 1 excitation per link), and $\omega_0 = 0.05J$. The strength of the spin-phonon coupling is varied: (a) $\alpha = 0.2J$, (b) $\alpha = 0.1J$ (c) $\alpha = 0.05J$, (d) $\alpha = 0.025J$. The vertical scale is arbitrary but is the same for the different values of α .

FIG. 5. Raman spectrum versus ω/J for the A_{1g} symmetry. These results correspond to 6 spins coupled to one phonon excitation and $\alpha = 0.1J$. (a)-(c) are all displayed on the same scale, (d)-(f) have been rescaled. $\omega_0 = 0.3$ for (a) and (d), $\omega_0 = 0.1$ for (b) and (e), and $\omega_0 = 0.025$ for (c) and (f).

FIG. 6. Raman spectra for the A_{1g} symmetry for 8 spins coupled to one phonon excitation and also scattering from an average over configurations of disorder. (a) $\alpha = 0.1J$, $\omega_0 = 0.05J$, and disorder chosen from a Gaussian with variance $\sigma = 0.2J$. (b) $\alpha = 0.3J$, $\omega_0 = 0.3J$, and disorder chosen from a Gaussian with variance $\sigma = 0.5J$.

FIG. 7. Raman scattering for the B_{1g} symmetry using (a) the bare Heisenberg Hamiltonian ($\alpha = 0$), (b) with dynamical phonons $\alpha = 0.1J$, $\omega_0 = 0.05J$, (c) with dynamical phonons $\alpha = 0.3J$, $\omega_0 = 0.3J$. The two bottom curves correspond to static Gaussian disorder of variance $\sigma = 0.2J$ (d), and $\sigma = 0.5J$ (e).

FIG. 8. Raman scattering from a 3×4 non-periodic lattice in the A_{1g} and B_{1g} geometries. The spin interactions J_{ij} have static Gaussian disorder of variance $\sigma = 0.5$